



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

if the British banks would agree to issue statistics that would bear comparison with those now published by the member banks of the Federal Reserve System. It is confessedly a forlorn hope. Hence a proper understanding of the sparse information now available is especially essential to American and other foreign investigators.

THE PROBABLE ERROR OF THE DIFFERENCE BETWEEN TWO DIFFERENCES

BY J. ARTHUR HARRIS

Given measures of two pairs of variables, w , x , and y , z , it is sometimes necessary to determine the probable error of the difference, $d_1 - d_2$, where $d_1 = x - w$, $d_2 = z - y$.

A formula has doubtless been given somewhere in the extensive literature, but I do not recall having seen it. Since it must involve the value of $r_{d_1 d_2}$ when there is any reason to suspect that the two differences may be correlated, a highly convenient method (which may be used when the correlations between the variables themselves are required and when variation and correlation are such that the ordinary theory of probable errors may be employed) is that based on the moments and product moments of the original variates, taken about 0 as origin. The formula for $r_{d_1 d_2}$ has already been suggested¹ and employed in the study of growth increments.²

The moments $\Sigma(w)$, $\Sigma(w^2)$, $\Sigma(x)$, $\Sigma(x^2)$, . . . , and the product moments $\Sigma(wx)$, $\Sigma(wy)$, $\Sigma(wz)$, $\Sigma(xy)$, . . . , are generally required for other purposes of the investigation.

The constants for the differences are given by well-known formulas:

$$\bar{d}_1 = \bar{x} - \bar{w},$$

$$\sigma_{d_1}^2 = [\Sigma(w^2) + \Sigma(x^2) - 2\Sigma(wx)]/N - \bar{d}_1^2,$$

and similarly for d_2 , where the bars denote means and the sigmas denote standard deviations.

The product moment from which $r_{d_1 d_2}$ may be deduced is given by

$$\Sigma(d_1 d_2) = \Sigma(wy) + \Sigma(xz) - \Sigma(xy) - \Sigma(wz).$$

The standard deviation of $d_1 - d_2$ may then be written

$$\sigma_{(d_1 - d_2)}^2 = \sigma_{d_1}^2 + \sigma_{d_2}^2 - 2r_{d_1 d_2} \sigma_{d_1} \sigma_{d_2}.$$

¹ Harris, J. Arthur, "Formulae for the determination of the correlation of size and of growth increments in the developing organisms." *Proc. Soc. Exp. Biol. Med.*, Vol. XVIII, pp. 4-5, 1921.

² Harris, J. Arthur, and Reed, H. S. "Inter-periodic correlation in the analysis of growth." *Biol. Bull.*, Vol. XL, pp. 243-258, 1921.